

CALCULUS OPTION (Q 8, PAPER 2)

LESSON NO. 3: MACLAURIN SERIES

2006

8 (a) Derive the Maclaurin series for $f(x) = e^x$ up to and including the term containing x^3 .

2005

8 (b) (i) Derive the Maclaurin series for $f(x) = \ln(1+x)$ up to and including the term containing x^3 .

(ii) Use those terms to find an approximation for $\ln \frac{11}{10}$.

(iii) Write down the general term of the series $f(x)$ and hence show that the series converges for $-1 < x < 1$.

2004

8 (b) $f(x) = f(0) + \frac{f'(0)x}{1!} + \frac{f''(0)x^2}{2!} + \frac{f'''(0)x^3}{3!} + \dots$ is the Maclaurin series.

(i) Derive the first five terms of the Maclaurin series for e^x .

(ii) Hence write down the first five terms of the Maclaurin series for e^{-x} and deduce the first three non-zero terms of the series for $\frac{e^x + e^{-x}}{2}$.

(iii) Write the general term of the series for $\frac{e^x + e^{-x}}{2}$ and use the Ratio Test to show that the series converges for all x .

2003

8 (b) $f(x) = f(0) + \frac{f'(0)x}{1!} + \frac{f''(0)x^2}{2!} + \frac{f'''(0)x^3}{3!} + \dots$ is the Maclaurin series.

(i) Derive the Maclaurin series for $f(x) = \log_e(1+x)$ up to and including the term containing x^4 .

(ii) Write down the general term and use the Ratio Test to show that the series converges for $-1 < x < 1$.

2002

8 (c) The Maclaurin series for $\tan^{-1} x$ is $x - \frac{x^3}{3} + \frac{x^5}{5} - \frac{x^7}{7} + \dots$. The series is convergent when $|x| < 1$.

- (i) Write down the first four terms in the series expansion for $\tan^{-1} \frac{1}{2}$.
- (ii) Use the fact that $\tan^{-1} \frac{1}{2} + \tan^{-1} \frac{1}{3} = \frac{\pi}{4}$ to derive a series expansion for π , giving the terms up to and including seventh powers.
- (iii) Use these terms to find an approximation for π . Give your answer correct to four places of decimal.

2001

8 (b) $f(x) = f(0) + \frac{f'(0)x}{1!} + \frac{f''(0)x^2}{2!} + \frac{f'''(0)x^3}{3!} + \dots$ is the Maclaurin series for $f(x)$.

- (i) Derive the Maclaurin series for $f(x) = \sin x$ up to and including the term containing x^7 .
- (ii) Write down the general term and use the Ratio Test to show that the series converges for all $x \in \mathbf{R}$.

ANSWERS

2006 8 (a) $1 + x + \frac{x^2}{2!} + \frac{x^3}{3!}$

2005 8 (b) (i) $x - \frac{1}{2}x^2 + \frac{1}{3}x^3$ (ii) $\frac{143}{1500}$ (iii) $u_n = (-1)^{n-1} \frac{x^n}{n}$

2004 8 (b) (i) $1 + \frac{x}{1!} + \frac{x^2}{2!} + \frac{x^3}{3!} + \frac{x^4}{4!}$ (ii) $1 + \frac{x^2}{2!} + \frac{x^4}{4!}$ (iii) $u_n = \frac{x^{2n-2}}{(2n-2)!}$

2003 8 (b) (i) $x - \frac{1}{2}x^2 + \frac{1}{3}x^3 - \frac{1}{4}x^4$ (ii) $u_n = (-1)^{n-1} \frac{x^n}{n}$

2002 8 (c) (i) $\frac{1}{2} - \frac{1}{24} + \frac{1}{160} - \frac{1}{896} + \dots$ (ii) $4[\frac{1}{2} - \frac{1}{24} + \frac{1}{160} - \frac{1}{896} + \frac{1}{3} - \frac{1}{81} + \frac{1}{1215} - \frac{1}{15309}]$
(iii) 3.1409

2001 8 (b) (i) $x - \frac{x^3}{3!} + \frac{x^5}{5!} - \frac{x^7}{7!} + \dots$ (ii) $u_n = (-1)^{n-1} \frac{x^{2n-1}}{(2n-1)!}$